

BLOOD-THINNER STOP/RESTART GUIDELINES

DO NOT DISCONTINUE UNTIL PRESCRIBING PHYSICIAN CLEARS

<u>BLOOD THINNER MEDICATION</u>	<u>WHEN TO STOP</u>	<u>WHEN TO RESTART</u>
Aspirin 81 mg	Ok to remain on therapy	
Aspirin 325 mg	6 days prior to procedure	24 hours post procedure
Goody Powder	6 days prior to procedure	24 hours post procedure
<u>NSAIDs</u>		
Diclofenac (Voltaren)	6 day prior to procedure	24 hours post procedure
Ketorolac (Toradol)	6 day prior to procedure	24 hours post procedure
Ibuprofen (Aleve Motrin Advil)	6 day prior to procedure	24 hours post procedure
Etodolac (Lodine)	6 days prior to procedure	24 hours post procedure
Indomethacin (Indocin)	6 days prior to procedure	24 hours post procedure
Naproxen (Naprosyn)	6 days prior to procedure	24 hours post procedure
Meloxicam (Mobic)	6 days prior to procedure	24 hours post procedure
Nabumetone (Relafen)	6 days prior to procedure	24 hours post procedure
Oxaprozin (Daypro)	6 days prior to procedure	24 hours post procedure
Piroxicam (Feldene)	6 days prior to procedure	24 hours post procedure
<u>Phosphodiesterase Inhibitors</u>		
Cilostazol (Pletal)	2 days prior to procedure	24 hours post procedure
Dipyridamole (Persantine)	2 days prior to procedure	24 hours post procedure
<u>Anticoagulants</u>		
Coumadin (Warfarin)	5 days prior/normal INR	24 hours post procedure
Acenocoumarol	3 days prior/normal INR	24 hours post procedure
IV Heparin	4 hours prior to procedure	2-24 hours post procedure
SubQ Heparin	8-10 hours prior	2 hours post procedure
Lovenox (once daily)	12 hours prior	24 hours post procedure
Lovenox (twice daily)	24 hours prior	24 hours post procedure
Fibrinolytic agents	48 hours prior	48 hours post procedure
Fondaparinux (Arixtra)	4 days prior	24 hours post procedure
Pentosan (Elmiron)	24 hours prior	24 hours post procedure
<u>P2Y12 Inhibitors</u>		
Clopidogrel (Plavix)	7 days prior	24 hours post procedure
Prasugrel (ESENTRO)	7-10 days prior	24 hours post procedure
Ticagrelor (Brilinta)	5 days prior	24 hours post procedure
<u>New anticoagulants</u>		
Dabigatran (Pradaxa)	4-5 days prior 6 days prior if impaired renal function	24 hours post procedure
Rivaroxaban (Xarelto)	3 days prior	24 hours post procedure
Apixaban (Eliquis)	3-5 days prior	24 hours post procedure
<u>Glycoprotein IIb/IIIa Inhibitors</u>		
Abciximab (ReoPro)	2-5 days prior	8-12 hours post procedure
Eptifibatide (Integrilin)	8-24 hours prior	8-12 hours post procedure
Tirofiban (Aggrastat)	8-24 hours prior	8-12 hours post procedure

THESE ARE GENERAL GUIDELINES. IF YOU HAVE A HISTORY OF STROKE, MECHANICAL MITRAL VALVE, PULMONARY EMBOLISM OR OTHER HIGH RISK FACTOR PLEASE CONSULT YOUR PROVIDER

Patient Signature: _____ **Date:** _____

Print Name of Patient: _____ **Date of Birth:** _____